

Statement about Investigation of the Death of Rachel Ferguson on 9th June 2001

I, Dr. Raymond Fulton, MB, FRCP London, was Medical Director of Altnagelvin Hospitals H&SS Trust at the time of Rachel Ferguson's death on 09/06/01. I was responsible for investigating the circumstances of her death within the hospital and to make recommendations for any action to prevent recurrence.

On 12/06/01 I set up a Critical Incident Enquiry involving all relevant clinical staff to establish the clinical facts. As a result of this six Action Points were agreed and circulated to all present on 13/06/01 (*see attachment 1*).

On 14/06/01 following Action Point 1 Dr Nesbitt, Clinical Director for Anaesthetics, wrote to me saying he had found that Solution 18 was currently used in several hospitals in Northern Ireland. He said he had reviewed the literature, which had convinced him that Solution 18 should not be used in surgical paediatric patients. He stated that henceforth Solution 18 would not be used in these circumstances in Altnagelvin (*see attachment 2*).

On 18/06/01 at a meeting of Medical Directors with Dr. I Carson, Medical Advisor to the CMO, at Castle Buildings I described the circumstances of this death. There were several anaesthetists present, some of whom said that they had heard of similar situations though it was not clear if there had been fatalities. I suggested that there should be regional guidelines.

On 22/07/01 I rang the Chief Medical Officer, Dr Henrietta Campbell, and informed her of the circumstances of the death. I suggested she should publicise the dangers of hyponatraemia when using low saline solutions in surgical children. I said there was a need for regional guidelines. Dr Campbell suggested that CREST (*Regional Guidelines Group*) might do this.

In Mid June 2001 I rang Dr. W McConnell, the Director of Public Health at the Western Health & Social Services Board, and described the circumstances of the death. He said he would discuss the matter at his next meeting with the Chief Medical Officer and the Directors of Public Health of the three other Health Boards. I sent him reprints from the British Medical Journal on Hyponatraemia (*see attachment 3*)

On 05/07/01 Dr McConnell wrote to confirm that he had discussed the case with the CMO and DPHs. Each DPH had agreed to alert the Paediatricians in their respective Board areas to the hazards of Hyponatraemia (*see attachment 4*).

On 6/07/01 Mrs Burnside, Chief Executive of Altnagelvin Hospitals H&SS Trust, contacted the CMO to personally advocate a regional review (*see attachment 5*). I remember seeing a reply from the CMO agreeing to set up a regional Enquiry Group and that Dr Nesbitt would be a member.

UTV

069A-087-339

On 14/01/02 I arranged for the CMO to view a presentation by Dr Nesbitt on Hyponatraemia while she was visiting Altnagelvin Hospital to present accreditation to the Trust's HSDU.

On 09/04/02 I chaired a meeting of relevant clinical staff to review the Action Plan of 12/06/01 in light of the publication of Departmental Guidelines on Hyponatraemia. (*see attachment 6*)

On 01/05/02 Dr. Nesbitt, who succeeded me as Medical Director, wrote to the CMO enquiring if the death of a child some years previously from hyponatraemia in the RBHSC had been reported to the Department. Dr. Nesbitt had become aware of the RBHSC case whilst investigating the death of Rachel Ferguson (*see attachment 7*).

On 10/05/02 Dr. Campbell replied that the Department had not been made aware of the first case either by the Royal Victoria Hospital or the Coroner. She had only become aware of the original case whilst working on developing guidelines following the death of Rachel Ferguson (*see attachment 8*).

Throughout this process I was struck by the wish of all concerned to learn from this death, which is unique in their experience. I received full co-operation from all clinical staff who are extremely distressed by Rachel's death.



DR RAYMOND FULTON
12/11/02

attachments 1-8

①

AGREED ACTION FOLLOWING CRITICAL INCIDENT MEETING 12/06/01

1 Review evidence for use of routine post-operative low electrolyte IV infusion and suggest changes if evidence indicates. No change in current use of Solution 18 until review.

Action Dr Nesbitt

2 Arrange daily U&E on all post-operative children receiving IV infusion on Ward 6.

Action Sister Miller

3 Inform surgical junior staff to assess these results promptly.

Action Mr Gilliland

4 All urinary output should be measured and recorded while IV infusion ~~progress~~ in progress.

Action Sister Miller

5 A chart for IV fluid infusion rates to be displayed on Ward 6 to guide junior medical staff.

Action Dr McCord

6 Review fluid balance documentation used on Ward 6.

Action A Witherow

R A FULTON
Medical Director

13/06/01

2

Dr G A Nesbitt
Clinical Director
Anaesthetic Department
Altnagelvin Hospital

Date: 14th June 2001

14 JUN 2001

Dr Raymond Fulton
Medical Director

Re: Fluid management in Children

Dear Dr Fulton, *Raymond.*

I have contacted several hospitals including The Royal Hospital for Sick Children and made enquiries about peri-operative fluid management.

The Children's Hospital Anaesthetists have recently changed their practise and have moved away from No.18 solution (fifth normal NaCl in 4% Dextrose) to Hartman's solution. This change occurred 6 months ago and followed several deaths involving No.18 solution.

Craigavon Hospital and the Ulster Hospital both use Hartman's intra-operatively and No.18 post-operatively as is our practise. The Anaesthetists in Craigavon have been trying to change the fluid regime to Hartman's postoperatively but have met resistance in the Paediatric wards where, as in Altnagelvin, they have followed a medical paediatric protocol.

In view of recent events, and papers on the subject, and the fact that the Children's Hospital no longer uses No.18 solution, I have decided to recommend that we do the same. I have spoken to Sister Miller in the Paediatric ward and also to Dr McCord who both are in agreement. Dr McCord has agreed to add this to the protocol he is developing for calculating the amount of fluid to be prescribed. He has further agreed that, pending discussion with his colleagues, fluid management in postoperative children should be under the supervision of paediatricians.

To summarise: Altnagelvin Hospital has followed what is a widespread and accepted policy of using No.18 solution for postoperative fluids. There is evidence to show that this policy is potentially unsafe in certain children who have undergone a surgical procedure. The Children's Hospital has ceased to use it and Craigavon is trying to effect a change in this direction. As from today we will no longer be routinely using this fluid in the management of surgical cases.

Yours sincerely,


G A Nesbitt Clinical Director

cc Theresa Brown

Risk Management Coordinator

UTV

069A-087-342

3

<input type="text"/>	<input type="text"/>
----------------------	----------------------

Home Help Search/Archive Feedback Search Result

BMJ 2001;322:780-782 (31 March)

Clinical review

Lesson of the week

Acute hyponatraemia in children admitted to hospital: retrospective analysis of factors contributing to its development and resolution

Do not infuse a hypotonic solution if the plasma sodium concentration is less than 138 mmol/l

<input type="checkbox"/> PDF of this article <input type="checkbox"/> Send a response to this article <input type="checkbox"/> Electronic responses to this article <input type="checkbox"/> PubMed citation <input type="checkbox"/> Related articles in PubMed <input type="checkbox"/> Download to Citation Manager <input type="checkbox"/> Search Medline for articles by: <input type="checkbox"/> Halberthal, M. Bohn, D. <input type="checkbox"/> Alert me when: <input type="checkbox"/> New articles cite this article
<input type="checkbox"/> Collections under which this article appears: <input type="checkbox"/> Other emergency medicine <input type="checkbox"/> Adolescents <input type="checkbox"/> Other Pediatrics

Michael Halberthal, fellow ^a, Mitchell L Halperin, professor ^b, Desmond Bohn, professor ^c.

^a Department of Critical Care Medicine, Hospital for Sick Children, Toronto, Ontario, Canada M5G 1X8, ^b Division of Nephrology, St Michael's Hospital Toronto, Toronto, Ontario, Canada M5B 1A6; ^c Department of Critical Care Medicine, Hospital for Sick Children, Toronto

Correspondence to: D J Bohn dbohn@sickkids.on.ca

Hyponatraemia (plasma sodium concentration less than 136 mmol/l) is acute if the decrease in natraemia occurs within 48 hours. The major dangers from this are brain cell swelling and herniation. ^{1 2} Two factors are required for hyponatraemia to develop: a source of electrolyte free water and vasopressin to prevent the excretion of that water. ³ Electrolyte free water is given routinely as maintenance fluids based on formulas developed in studies in healthy children more than 40 years ago. ^{4 5} There are many reasons to anticipate that vasopressin will be released in sick patients (box). ⁶ Patients with an acute illness may arrive in hospital with a low plasma sodium concentration because of previous water intake. Hence, to minimise the potential threat of brainstem herniation it is important to measure the plasma sodium concentration if intravenous solutions are to be given.

UTV

Causes of vasopressin release

- Hypernatraemia (most important stimulus, but not in these patients)
- Low "effective" circulating volume (greater than 7% decrease in extracellular fluid volume)
- Nausea, pain, anxiety
- Drugs (some act through inducing nausea)
- Afferent stimuli by way of the vagus nerve for example, lung lesions

069A-087-343

- Disturbances of the central nervous system (meningitis, encephalitis)
- Metabolic and endocrine disorders for example, hypothyroidism, hypo adrenalism, porphyria

We describe symptomatic hyponatraemia developing over 48 hours in children. In each patient, hypotonic solutions were infused using current guidelines.⁷ We related the volume of electrolyte free water given to the decrease in natraemia and assessed whether actions of vasopressin persisted to guide emergency corrective therapy.⁸

We reviewed all patient charts (306 charts) with a recorded diagnosis of hyponatraemia for the past 10 years. Patients were included if their decrease in natraemia was to less than 130 mmol/l and this occurred within 48 hours, if intravenous fluids were given, and if an underlying disease did not compromise renal handling of sodium or water. Thirty patients had acute hyponatraemia. Crucial information was missing for seven, leaving 23 patients in the study group. The median age was five years (range one month to 21 years), with males predominating (18 of 23); 13 developed hyponatraemia in the postoperative period. Fifteen patients were referred to the critical care unit after the development of symptomatic hyponatraemia while receiving intravenous fluids. 11 were from the hospital wards and four were transferred from other institutions. Symptoms included seizures (18 patients) and vomiting, 17 a warning sign of an increased intracranial pressure. Treatment was withdrawn from five patients after brainstem coning. One patient sustained permanent, severe neurological damage.



Results

All the children received hypotonic fluids while their plasma sodium concentration was less than 140 mmol/l, because of the wide belief in paediatric practice that "maintenance fluids" should be hypotonic.⁹ In fact the volume of maintenance fluid given was 50% greater than recommended values in 16 of the 23 patients. This infusion of hypotonic fluids increased the risk of acute hyponatraemia and brain swelling because vasopressin is typically present in this setting.^{1 2 10 11} In quantitative terms, some of the electrolyte free water infused was retained in six of the patients because their urine sodium plus potassium concentration was less than 25 mmol/l (fig 1). In six patients more electrolyte free water was infused than needed to cause the observed decline in natraemia (points above line of identity in fig 2). The remainder of the patients had a decrease in natraemia that exceeded the decline if the entire volume of electrolyte free water infused was retained (points below broken line in fig 2). Therefore there was either another non-recorded input of water or the excretion of a large volume of hypertonic urine (a desalination of infused isotonic saline¹²).

<input type="checkbox"/>	Top
<input type="checkbox"/>	Results
<input type="checkbox"/>	Discussion
<input type="checkbox"/>	References

UTV

069A-087-344



Discussion

UTV

<input type="checkbox"/>	Top
<input type="checkbox"/>	Results
<input type="checkbox"/>	Discussion
<input type="checkbox"/>	References

One objective of our study was to assess the renal actions of vasopressin. Because six patients had very hypotonic urine at their recorded nadirs of natriuria, their plasma sodium concentration might have been much lower before water diuresis began (fig 1). Had their plasma sodium concentration been measured after this large water diuresis, the erroneous conclusion might have been drawn that acute hyponatraemia had never been present. Hence its incidence may be much higher than shown by an analysis of hospital records. Therefore acute hyponatraemia could have been an occult cause of morbidity and mortality.

Another implication of cessation of the release of vasopressin concerns treatment. Treatment for acute, symptomatic hyponatraemia causes a prompt decline in the size of brain cells.¹⁰ Hypertonic saline (3%) is the commonest treatment for shrinking brain cell volume, thereby lowering intracranial pressure. Treatment must be prompt because deterioration may be rapid and irreversible, even when symptoms are mild. Enough hypertonic saline (a total of 5 mmol of

069A-087-345

sodium chloride per litre of body water¹³) is needed acutely to lower intracranial pressure sufficiently to minimise this risk (the plasma sodium concentration should be increased by 5 mmol/l over several hours). Because an excessively rapid rate of correction of hyponatraemia might have deleterious effects,⁶ hypertonic saline should not be given if there is a brisk water diuresis. For example, the plasma sodium concentration will also increase by 1.2 mmol/l/h if 6 ml of electrolyte free water are excreted per kilogram per hour (total body water is close to 600 ml/kg; 6 ml is a 1% change of 120 mmol/l). Whereas excretion of hypotonic urine indicates that electrolyte free water is being excreted (6 of 17 patients, fig 1), it is also important to consider the rate of urine flow. Little electrolyte free water was excreted in the index oliguric patient (flow 0.16 ml/kg/h). By contrast, the excretion of electrolyte free water was high enough to increase the plasma sodium concentration by close to 3 mmol/l/h in the polyuric index patient who recovered (15 ml/kg/h). Vasopressin continued to act in patients excreting isotonic or hypertonic urine, so hypotonic intake must be avoided in them. With these high urine tonicities a further decrease in natraemia would be anticipated if the urine output was high (index case designated with a urine output of 5.3 ml/kg body weight, fig 1).¹² Finally, vasopressin concentrations may decline abruptly, increasing the excretion of electrolyte free water.

Serious symptoms may become evident when hyponatraemia approaches 120 mmol/l, but there are cases where symptoms become evident with a higher plasma sodium concentration, whereas others tolerate this electrolyte disorder without developing seizures.¹⁴ Apart from underlying conditions that might make a patient more susceptible to seizures, a possible important factor could be the extracellular fluid volume of the brain. If this volume was expanded by a large infusion of isotonic saline, a higher intracranial pressure might be present at a given degree of hyponatraemia. Moreover, because there is a relatively larger proportion of brain cell volume to extracellular fluid volume in young patients, they are more vulnerable to an increase in brain cell volume.

Study limitations

Because of a reporting and referral bias, the incidence of adverse outcomes from hyponaetremia cannot be deduced from these data. Our results highlight the dangers of the routine use of hypotonic solutions when vasopressin acts. The currently used guidelines for maintenance fluids in children admitted to hospital must be changed because they do not take into account the unpredictability of vasopressin secretion. We recommend that the concentration of plasma sodium should be measured when starting an intravenous infusion. If it is less than 140 mmol/l then isotonic and not hypotonic fluids should be given. The use of hypotonic solutions should be reserved for patients who have a plasma sodium concentration greater than 140 mmol/l. If a patient receives intravenous fluid that exceeds 5% of total body water (30 ml/kg) then their plasma sodium concentration should be measured. If an intravenous infusion is started to give drugs, a small volume should be used, and the solution should be isotonic if possible.

□ Acknowledgments

UTV

069A-087-346

Contributors: MH collected the data and drafted the original manuscript. MLH analysed the data and coauthored the manuscript. DB had the original idea and coauthored the manuscript; he will act as guarantor for the paper.

Footnotes

Competing interests: None declared.

References

<input type="checkbox"/> Top
<input type="checkbox"/> Results
<input type="checkbox"/> Discussion
<input type="checkbox"/> References

1. Arieff AI. Hyponatremia, convulsions, respiratory arrest, and permanent brain damage after elective surgery in healthy women. *N Engl J Med* 1986; 314: 1529-1535[[Abstract](#)].
2. Arieff AI, Ayus JC, Fraser CL. Hyponatremia and death or permanent brain damage in healthy children. *BMJ* 1992; 304: 1218-1222[[Medline](#)].
3. Halperin ML, Oh MS. The dysnatremias: hyponatremia and hypernatremia. In: Glasscock RJ, ed. *Current therapy in nephrology and hypertension*. 4th ed. Philadelphia: Mosby-Year Book, 1998:1-9.
4. Darrow DC, Pratt EL. Fluid therapy. *JAMA* 1950; 143: 365-373.
5. Halliday M, Segar W. The maintenance need for water in parenteral fluid therapy. *Pediatrics* 1957; 19: 823-832.
6. Sterns RH, Clark EC, Silver SM. Clinical consequences of hyponatremia and its correction. In: Seldin DW, Giebisch G, eds. *Clinical disorders of water metabolism*. New York: Raven, 1993:225-236.
7. Adelman RD, Solhung MJ. Pathophysiology of body fluids and fluid therapy. In: Nelson WE, ed. *Textbook of pediatrics*. 15th ed. Philadelphia: WB Saunders, 1996:207.
8. Rose BD. New approach to disturbances in the plasma sodium concentration. *Am J Med* 1986; 81: 1033-1040[[Medline](#)].
9. Chesney RW. The maintenance need for water in parenteral fluid therapy. *Pediatrics* 1998; 102: 399-400[[Full Text](#)].
10. Soupart A, Decaux G. Therapeutic recommendations for management of severe hyponatremia: current concepts on pathogenesis and prevention of neurologic complications. *Clin Nephrol* 1996; 46: 149-169[[Medline](#)].
11. Wattad A, Chiang M, Leighton-Hill L. Hyponatremia in hospitalized children. *Clin Pediatr (Phila)* 1992; 31: 153-157[[Medline](#)].
12. Steele A, Gowrishankar M, Abrahamson S, Mazer D, Feldman R, Halperin ML. Postoperative hyponatremia despite isotonic saline infusion: a phenomenon of 'desalination.' *Ann Intern Med* 1997;126:20-5.
13. Spital A, Sterns RD. The paradox of sodium's volume of distribution: why an extracellular solute appears to distribute over total body water. *Arch Intern Med* 1989; 149: 1255-1257 [[Medline](#)].
14. Ayus JC, Wheeler JM, Arieff AI. Postoperative hyponatremic encephalopathy in menstruant women. *Ann Intern Med* 1992; 117: 891-897[[Medline](#)].

UTV

(Accepted 26 July 2000)


© BMJ 2001

069A-087-347

43

WESTERN
HEALTH AND SOCIAL SERVICES BOARD

AREA BOARD HEADQUARTERS



WMcC/HP

5 July 2001

Dr R Fulton
Medical Director
Altnagelvin Health & Social Services Trust
Glenshane Road
Altnagelvin
Londonderry

Dear Raymond

As we agreed, I raised this issue for discussion at the recent CMO/DPH meeting. Dr Campbell indicated that her CMO update was already at the printers and therefore it would be difficult to include anything within it. Knowing this, it was suggested that I would write to each of my Director of Public Health colleagues who would then take responsibility for drawing the issue to the attention of any relevant paediatric settings within their respective Boards.

I enclose a copy of the letter which I have sent to them and I have also taken the liberty of copying to them the articles which you helpfully sent to me. I hope you do not mind my suggestion within the letter that anyone who wanted more detailed discussion might contact you for further information.

Best wishes.

Yours sincerely

Dr W W M McConnell
DIRECTOR OF PUBLIC HEALTH

Enc

UTV

069A-087-348

42

WMcC/HP

5 July 2001

To:

Directors of Public Health

Dear

I am writing to you, as agreed at our recent meeting, with further details regarding the unfortunate death of a young girl at Altnagelvin Hospital. It appears that the use of hypotonic saline is still common practice in a number of paediatric units although there has been information around for a few years suggesting this does present risks to a very small number of children in the acute perioperative period. I have enclosed some articles which you might find of benefit in relation to this.

I know from discussions about this issue with Dr Raymond Fulton, Medical Director at Altnagelvin Hospital, that some paediatric settings within Northern Ireland have made appropriate changes but that in others, while the information may be known by anaesthetic staff, there has not necessarily been discussion regarding change between anaesthetists, surgeons and paediatricians.

I felt it might be helpful to draw this issue to your attention in order that each of us can check with those paediatric settings in our respective Boards to see whether modification in practice may be needed. I hope that the articles are helpful to you. If further more specific information is required I am sure that Dr Fulton would be happy to discuss this with anyone who contacted him.

Best wishes.

Yours sincerely

Dr W W M McConnell
DIRECTOR OF PUBLIC HEALTH

Enc

UTV

069A-087-349

(4) (5)

Sally Doherty

From: Sally Doherty
Sent: 26 July 2001 11:08
To: 'Henrietta.campbell@' [REDACTED]
Subject: electrolyte balance in post operative children

I am writing further to Dr Fulton's conversation with you regarding the above, and am concerned to ensure that an overview of the research evidence is being undertaken. I believe that this is a regional, as opposed to local hospital issue, and would emphasise the need for a critical review of evidence.

I would be extremely grateful if you would ensure that the whole of the medical fraternity learned of the shared lesson.

I await to hear further from you.

Stella.

UTV

069A-087-350

CRITICAL INCIDENT REVIEW MEETING 09/4/02

To review the Action Plan of the critical incident meeting of 12-6-2001 following the death of Rachel Ferguson.

- 1 Review evidence for use of routine post-operative low electrolyte IV infusion and suggest changes if evidence indicates. No change in current use of Solution 18 until review.
 - An immediate review was undertaken and a decision was taken that from all Surgical patients (including orthopaedic) to receive I V Hartmans Solution and 6 hourly BM's.
- 2 Arrange daily U&E on all post-operative children receiving IV infusion on Ward 6.
 - This was immediately actioned by Sister Miller. The phlebotomists take the blood. It is not clear who is responsible for ordering the blood. Mrs Witherow and Mrs Brown will prepare ward guidelines

Acion T Brown A Witherow

- 3 Inform surgical junior staff to assess these results promptly.
 - This was immediately actioned by Mr Gilliland. All staff are informed at induction. This information should be included in the Junior Doctors handbook. At the moment blood results come up on the computer. This does not show the normal range. Agreed that all bloods are to be reported to the Surgeons routinely. Anne Witherow to speak to Dr. M O'Kane to ascertain if the normal ranges can be put on the computer.

Action A Witherow

- 4. All urinary output should be measured and recorded while IV infusion is in progress.
 - The fluid balance sheet has been revised to allow recording of urinary output and vomit.

5. A chart for IV fluid infusion rates to be displayed on Ward 6 to guide junior medical staff.
 - The chart was prepared and displayed by Dr Mc Cord by July 2001.
6. Review fluid balance documentation used on Ward 6.
 - The fluid balance sheet has been revised to show exact timing of IV Fluids, and when they have been discontinued. It was noted that there is a Regional Group currently reviewing this form. We will await receipt of the revised form.
7. Need to agree responsibility for the prescribing and management of fluids post operatively. Agreed that Dr. Nesbitt will discuss with Anaesthetists and agree a maximum time that postoperative fluids will be prescribed by anaesthetists.

Action Dr. Nesbitt

8. Departmental guidelines received April 2002 regarding fluid management in all children have been displayed on ward 6, theatres and A&E.



R A FULTON
11-4-2002

7

Dr G A Nesbitt
Medical director
Altnagelvin Hospital

1st May 2002

Dr Henrietta Campbell
Chief Medical Officer
Castle Buildings
Upper Newtownards Road
Belfast BT4 3SJ

Re: Hyponatraemia in Children receiving intravenous fluids

Dear Dr Campbell,

Following the death of a child in Altnagelvin Hospital, which is thought to have followed severe hyponatraemia, many steps have been taken to ensure that such an event does not occur again. We are all anxious to learn from what was a dreadful experience and to share vital information with others. Guidance issued from your Department will help in this regard and we are grateful for the recent posters on the subject.

I am interested to know if any such guidance was issued by the Department of Health following the death of a child in the Belfast Hospital for Sick Children which occurred some 5 years ago and whose death the Belfast Coroner investigated. I was unaware of this case and am somewhat at a loss to explain why.

I would be grateful if you could furnish me with any details of that particular case for I believe that questions will be asked as to why we did not learn from what appears to have been a similar event.

Yours sincerely,

G A Nesbitt Medical Director

UTV

069A-087-353

8

Department of Health, Social Services & Public Safety
An Roinn Sláinte, Seirbhísí Sóisialta agus Sábháilteachta Poiblí

From The Chief Medical Officer:
Dr Henrietta Campbell CB


Castle Buildings
Upper Newtownards Road
Belfast BT4 3SJ

Telephone: [REDACTED]
Fax: [REDACTED]

E-Mail: [REDACTED]

Dr G A Nesbitt
Medical Director
Altnagelvin Hospitals HSS Trust
Altnagelvin Area Hospital
Glenshane Road
LONDONDERRY
BT47 6SB

10 May 2002



Dear Dr Nesbitt

HYPONATRAEMIA IN CHILDREN RECEIVING INTRAVENOUS FLUIDS

Thank you for your letter of 1 May regarding guidelines for the prevention of hyponatraemia in children receiving intravenous fluids.

Your letter referred to a Coroner's case five years ago in which the cause of death of a child was reported to be due to hyponatraemia. This Department was not made aware of the case at the time either by the Royal Victoria Hospital or the Coroner. We only became aware of that particular case when we began the work of developing guidelines following the death at Altnagelvin.

I would like to thank the staff at Altnagelvin for alerting me to the need for guidelines and for all the assistance which was given to their development.

With kind regards.

Yours sincerely



HENRIETTA CAMPBELL (Dr)

069A-087-354

UTV

Copy to
[Handwritten notes]



Department of Health, Social Services & Public Safety
An Roinn Sláinte, Seirbhísí Sóisialta agus Sábháilteachta Poiblí

From The Chief Medical Officer:
Dr Henrietta Campbell CB

Castle Buildings
Upper Newtownards Road
Belfast BT4 3SJ

Telephone: [REDACTED]

Fax: [REDACTED]

E-Mail: [REDACTED]

Medical Directors of Acute Trusts
Directors of Nursing in Acute Trusts
Consultant Paediatricians
Consultant Surgeons
Consultant Neurosurgeons
Consultant Anaesthetists/Intensivists
Consultants in Plastic Surgery/Burns
Consultants in A&E Medicine
Consultant Pathologists

25 March 2002

Dear Colleagues

PREVENTION OF HYPONATRAEMIA IN CHILDREN

Guidance on the Prevention of Hyponatraemia in Children has been published and will be forwarded to you under separate cover. It has been prepared as an A2 sized poster and I ask you to ensure that the posters are prominently displayed in all units that may accommodate children. The Guidance has been developed by a multidisciplinary working group established by the Department and the work has been supported and endorsed by CREST.

Hyponatraemia can be extremely serious and has in the past few years been responsible for two deaths among children in Northern Ireland. Hyponatraemia is a problem of water balance and most often reflects the failure to excrete water. Stress, pain and nausea are all potential stimulators of the antidiuretic hormone ADH which inhibits water excretion.

Any child receiving IV fluids or oral rehydration is potentially at risk of hyponatraemia. The administration of excess or inappropriate fluid to a sick child may result in serious or life threatening hyponatraemia. There is a particular concern about the use of 0.18% Sodium Chloride in Glucose among children as it has been implicated in cases of hyponatraemia. While it may pose a risk because of the relatively low sodium content no specific fluid is without risk. This has been emphasised in a recent letter received from the Medicines Control Agency which stated that while hyponatraemia is a risk with 0.18% Sodium Chloride, electrolyte imbalance is a risk with all intravenous solutions.

The Guidance emphasises that every child receiving intravenous fluids requires a thorough baseline assessment, that fluid requirements must be calculated accurately and fluid balance must be rigorously monitored. Following this simple advice will prevent children from developing hyponatraemia.

The Guidance is designed to provide general advice and does not specify particular fluid choices. Fluid protocols should be developed locally to complement the Guidance and provide more specific direction to junior staff. This is particularly important in subspecialty areas such as renal medicine, burns units and

UTV

069A-087-355



neurosurgery. It will be important to audit compliance with the guidance and locally developed protocols and to learn from clinical experiences.

I would like to extend my thanks to all members of the multidisciplinary group who have worked together to provide clear and practical guidance to improve the care of sick children. The Guidance is also available on the Departmental website www.dhsspsni.gov.uk.

Yours sincerely



HENRIETTA CAMPBELL (Dr)