

OPERATION NOTES

Affix label or enter

Adam Strain 364377 Musg. Wd.

Operation performed Insertion Central Line, Cystoscopy & Retrograde 29/5/92

Surgeon Mr W A McCallion / Mr R J Stewart / Mr S Brown Anaesthetist Dr McCarthy / Crean

Assistant Sister Malcolmson

Incision

Indication: Pre base bilateral ureteric reimplant. Left ureter sloughed?

Findings requiring left to right TUO now presents with recurring UTI's

query due to obstructive uropathy . Requires long term antibiotics.

1) Insertion Broviac Line into left common facial vein. Transverse cervical incision. Left common facial vein identified, entering left internal

Procedure jugular. Left common facial ligated with 5 X 0 PDS. Broviac line tunnelled

from anterior chest wall using Westminster and inserted into common facial

vein and then internal jugular. Check X-ray confirmed tip of broviac line

in proximal SBC. Neck wound closed in layers of 5 X 0 PDs and wound anterior

chest wall closed 5 X 0 PDS.

2) Cystoscopy using 30° telescope and 11 storz sheath. The right ureteric orifice was identified in the mid line approx at the level of inter ureteric

bar. No lesion was found within the bladder. The UO was easily catheterised

with a size 3 ureteric catheter. The cystoscope was withdrawn and the baby

returned to X-ray for retrograde pyelogram.

W A McCallion

CMcI

Blood

Closure

Drains

Royal Belfast Hospital for Sick Children
Belfast, BT12 6BE