



FOR SMOKING CESSATION
Z MARKS THE SPOT

Zyban
bupropion HCl SR
Science against smoking

[Home](#) [Help](#) [Search/Archive](#)

[Feedback](#)

[Search Result](#)

BMJ 2001;322:780-782 (31 March)

Clinical review

Lesson of the week

Acute hyponatraemia in children admitted to hospital: retrospective analysis of factors contributing to its development and resolution

Do not infuse a hypotonic solution if the plasma sodium concentration is less than 138 mmol/l

[PDF of this article](#)

[Send a response to this article](#)

[Electronic responses to this article](#)

[PubMed citation](#)

[Related articles in PubMed](#)

[Download to Citation Manager](#)

[Search Medline for articles by:](#)

Halberthal, M. || Bohn, D.

[Alert me when:](#)

[New articles cite this article](#)

[Collections under which this article appears:](#)

[Other emergency medicine](#)

[Adolescents](#)

[Other Pediatrics](#)

Michael Halberthal, fellow ^a, Mitchell L Halperin, professor ^b, Desmond Bohn, professor ^c

^a Department of Critical Care Medicine, Hospital for Sick Children, Toronto, Ontario, Canada M5G 1X8, ^b Division of Nephrology, St Michael's Hospital Toronto, Toronto, Ontario, Canada M5B 1A6, ^c Department of Critical Care Medicine, Hospital for Sick Children, Toronto

Correspondence to: D J Bohn dbohn@sickkids.on.ca

Hyponatraemia (plasma sodium concentration less than 136 mmol/l) is acute if the decrease in natraemia occurs within 48 hours. The major dangers from this are brain cell swelling and herniation.^{1 2} Two factors are required for hyponatraemia to develop: a source of electrolyte free water and vasopressin to prevent the excretion of that water.³ Electrolyte free water is given routinely as maintenance fluids based on formulas developed in studies in healthy children more than 40 years ago.^{4 5} There are many reasons to anticipate that vasopressin will be released in sick patients (box).⁶ Patients with an acute illness may arrive in hospital with a low plasma sodium concentration because of previous water intake. Hence, to minimise the potential threat of brainstem herniation it is important to measure the plasma sodium concentration if intravenous solutions are to be given.

Causes of vasopressin release

- Hypernatraemia (most important stimulus, but not in these patients)
- Low "effective" circulating volume (greater than 7% decrease in extracellular fluid volume)
- Nausea, pain, anxiety
- Drugs (some act through inducing nausea)
- Afferent stimuli by way of the vagus nerve—for example, lung lesions
- Disturbances of the central nervous system (meningitis, encephalitis)
- Metabolic and endocrine disorders—for example, hypothyroidism, hypoadrenalism, porphyria

RF – ROYAL

064-021-058


We describe symptomatic hyponatraemia developing over 48 hours in children. In each patient, hypotonic solutions were infused using current guidelines.⁷ We related the volume of electrolyte free water given to the decrease in natraemia and assessed whether actions of vasopressin persisted to guide emergency corrective therapy.⁸

We reviewed all patient charts (306 charts) with a recorded diagnosis of hyponatraemia for the past 10 years. Patients were included if their decrease in natraemia was to less than 130 mmol/l and this occurred within 48 hours, if intravenous fluids were given, and if an underlying disease did not compromise renal handling of sodium or water. Thirty patients had acute hyponatraemia. Crucial information was missing for seven, leaving 23 patients in the study group. The median age was five years (range one month to 21 years), with males predominating (18 of 23); 13 developed hyponatraemia in the postoperative period. Fifteen patients were referred to the critical care unit after the development of symptomatic hyponatraemia while receiving intravenous fluids—11 were from the hospital wards and four were transferred from other institutions. Symptoms included seizures (18 patients) and vomiting, 17 a warning sign of an increased intracranial pressure. Treatment was withdrawn from five patients after brainstem coning. One patient sustained permanent, severe neurological damage.

Results

All the children received hypotonic fluids while their plasma sodium concentration was less than 140 mmol/l, because of the wide belief in paediatric practice that "maintenance fluids" should be hypotonic.⁹ In fact the volume of maintenance fluid given was 50% greater than recommended values in 16 of the 23 patients. This infusion of hypotonic fluids increased the risk of acute hyponatraemia and brain swelling because vasopressin is typically present in this setting.^{1 2 10 11} In quantitative terms, some of the electrolyte free water infused was retained in six of the patients because their urine sodium plus potassium concentration was less than 25 mmol/l (fig 1). In six patients more electrolyte free water was infused than needed to cause the observed decline in natraemia (points above line of identity in fig 2). The remainder of the patients had a decrease in natraemia that exceeded the decline if the entire volume of electrolyte free water infused was retained (points below broken line in fig 2). Therefore there was either another non-recorded input of water or the excretion of a large volume of hypertonic urine (a desalination of infused isotonic saline¹²).

Top
• Results
Discussion
References



RF - ROYAL

[in this window]
[in a new window]




Fig 2. Comparison of decline in plasma sodium concentration with amount of electrolyte free water given in 17 patients. Difference in sodium concentration was between initial value and that at its nadir

View larger version (26K):
[in this window]
[in a new window]

Discussion

One objective of our study was to assess the renal actions of vasopressin. Because six patients had very hypotonic urine at their recorded nadirs of natraemia, their plasma sodium concentration might have been much lower before water diuresis began (fig 1). Had their plasma sodium concentration been measured after this large water diuresis, the erroneous conclusion might have been drawn that acute hyponatraemia had never been present. Hence its incidence may be much higher than shown by an analysis of hospital records. Therefore acute hyponatraemia could have been an occult cause of morbidity and mortality.

Top
Results
• Discussion
References

Another implication of cessation of the release of vasopressin concerns treatment. Treatment for acute, symptomatic hyponatraemia causes a prompt decline in the size of brain cells.¹⁰ Hypertonic saline (3%) is the commonest treatment for shrinking brain cell volume, thereby lowering intracranial pressure. Treatment must be prompt because deterioration may be rapid and irreversible, even when symptoms are mild. Enough hypertonic saline (a total of 5 mmol of sodium chloride per litre of body water¹³) is needed acutely to lower intracranial pressure sufficiently to minimise this risk (the plasma sodium concentration should be increased by 5 mmol/l over several hours). Because an excessively rapid rate of correction of hyponatraemia might have deleterious effects,⁶ hypertonic saline should not be given if there is a brisk water diuresis. For example, the plasma sodium concentration will also increase by 1.2 mmol/l/h if 6 ml of electrolyte free water are excreted per kilogram per hour (total body water is close to 600 ml/kg; 6 ml is a 1% change of 120 mmol/l). Whereas excretion of hypotonic urine indicates that electrolyte free water is being excreted (6 of 17 patients, fig 1), it is also important to consider the rate of urine flow. Little electrolyte free water was excreted in the index oliguric patient (flow 0.16 ml/kg/h). By contrast, the excretion of electrolyte free water was high enough to increase the plasma sodium concentration by close to 3 mmol/l/h in the polyuric index patient who recovered (15 ml/kg/h). Vasopressin continued to act in patients excreting isotonic or hypertonic urine, so

hypotonic intake must be avoided in them. With these high urine tonicities a further decrease in hyponatraemia would be anticipated if the urine output was high (index case designated with a urine output of 5.3 ml/kg body weight, fig 1).¹² Finally, vasopressin concentrations may decline abruptly, increasing the excretion of electrolyte free water.

Serious symptoms may become evident when hyponatraemia approaches 120 mmol/l, but there are cases where symptoms become evident with a higher plasma sodium concentration, whereas others tolerate this electrolyte disorder without developing seizures.¹⁴ Apart from underlying conditions that might make a patient more susceptible to seizures, a possible important factor could be the extracellular fluid volume of the brain. If this volume was expanded by a large infusion of isotonic saline, a higher intracranial pressure might be present at a given degree of hyponatraemia. Moreover, because there is a relatively larger proportion of brain cell volume to extracellular fluid volume in young patients, they are more vulnerable to an increase in brain cell volume.

Study limitations

Because of a reporting and referral bias, the incidence of adverse outcomes from hyponatraemia cannot be deduced from these data. Our results highlight the dangers of the routine use of hypotonic solutions when vasopressin acts. The currently used guidelines for maintenance fluids in children admitted to hospital must be changed because they do not take into account the unpredictability of vasopressin secretion. We recommend that the concentration of plasma sodium should be measured when starting an intravenous infusion. If it is less than 140 mmol/l then isotonic and not hypotonic fluids should be given. The use of hypotonic solutions should be reserved for patients who have a plasma sodium concentration greater than 140 mmol/l. If a patient receives intravenous fluid that exceeds 5% of total body water (30 ml/kg) then their plasma sodium concentration should be measured. If an intravenous infusion is started to give drugs, a small volume should be used, and the solution should be isotonic if possible.

Acknowledgments

Contributors: MH collected the data and drafted the original manuscript. MLH analysed the data and coauthored the manuscript. DB had the original idea and coauthored the manuscript; he will act as guarantor for the paper.

Footnotes

Competing interests: None declared.

References

[Top](#)
[Results](#)
[Discussion](#)

• [References](#)

1. Arieff AI. Hyponatremia, convulsions, respiratory arrest, and permanent brain damage after elective surgery in healthy women. *N Engl J Med* 1986; 314: 1529-1535[Abstract].
2. Arieff AI, Ayus JC, Fraser CL. Hyponatremia and death or permanent brain damage in healthy children. *BMJ* 1992; 304: 1218-1222[Medline].
3. Halperin ML, Oh MS. The dysnatremias: hyponatremia and hypernatremia. In: Glascock RJ, ed. *Current therapy in nephrology and hypertension*. 4th ed. Philadelphia: Mosby-Year Book, 1998:1-9.
4. Darrow DC, Pratt EL. Fluid therapy. *JAMA* 1950; 143: 365-373.
5. Halliday M, Segar W. The maintenance need for water in parenteral fluid therapy. *Pediatrics* 1957; 19: 823-832.
6. Sterns RH, Clark EC, Silver SM. Clinical consequences of hyponatremia and its correction. In: Seldin DW, Giebisch G, eds. *Clinical disorders of water metabolism*. New York: Raven, 1993:225-236.
7. Adelman RD, Solhung MJ. Pathophysiology of body fluids and fluid therapy. In: Nelson WE, ed. *Textbook of pediatrics*. 15th ed. Philadelphia: WB Saunders, 1996:207.
8. Rose BD. New approach to disturbances in the plasma sodium concentration. *Am J Med* 1986; 81: 1033-1040[Medline].
9. Chesney RW. The maintenance need for water in parenteral fluid therapy. *Pediatrics* 1998; 102: 399-400[Full Text].
10. Soupart A, Decaux G. Therapeutic recommendations for management of severe hyponatremia: current concepts on pathogenesis and prevention of neurologic complications. *Clin Nephrol* 1996; 46: 149-169[Medline].
11. Wattad A, Chiang M, Leighton-Hill L. Hyponatremia in hospitalized children. *Clin Pediatr (Phila)* 1992; 31: 153-157[Medline].
12. Steele A, Gowrishankar M, Abrahamson S, Mazer D, Feldman R, Halperin ML. Postoperative hyponatremia despite isotonic saline infusion: a phenomenon of 'desalination.' *Ann Intern Med* 1997;126:20-5.
13. Spital A, Sterns RD. The paradox of sodium's volume of distribution: why an extracellular solute appears to distribute over total body water. *Arch Intern Med* 1989; 149: 1255-1257 [Medline].
14. Ayus JC, Wheeler JM, Arieff AI. Postoperative hyponatremic encephalopathy in menstruant women. *Ann Intern Med* 1992; 117: 891-897[Medline].

(Accepted 26 July 2000)

© BMJ 2001

Rapid Response responses to this article:

Read all Rapid Response responses

Hyponatraemia: nadirs and paradox of the missing volumetric overload.

Ahmed N. Ghanem, Consultant Urologist , King Khalid hospital, Najran, Saudi Arabia
bmj.com, 23 Apr 2001 [Response]

PDF of this article

Send a response to this article

Electronic responses to this article

PubMed citation

Related articles in PubMed

Download to Citation Manager

Search Medline for articles by:

Halberthal, M. || Bohn, D.

Alert me when:

New articles cite this article

Collections under which this article appears:

Other emergency medicine

Adolescents

Other Pediatrics