

## Editorial

# Postoperative hyponatraemic encephalopathy following elective surgery in children

ALLEN L. ARIEFF MD

Department of Medicine, University of California School of Medicine,  
San Francisco, CA, USA

## Introduction

In the United States, there are an estimated 15 000 deaths per year as a consequence of postoperative hyponatraemia (1) (Figure 1). There have been a number of recent studies which have described postoperative hyponatraemic encephalopathy with death or permanent brain damage (2-6). From these studies, it appears that brain damage associated with postoperative hyponatraemic encephalopathy primarily affects menstruant women (1) and prepubertal children (6).

### *Postoperative hyponatraemic encephalopathy in prepubertal children*

There are multiple reports of prepubertal children suffering brain damage from postoperative hyponatraemic encephalopathy (6-9). The aetiology of the hyponatraemia usually involves a combination of: a) intravenous hyponatraemic fluids; b) elevated plasma antidiuretic hormone (ADH); c) respiratory insufficiency secondary to hyponatraemic encephalopathy. It has been demonstrated in several series that plasma levels of ADH (vasopressin, antidiuretic hormone) are elevated in virtually every postoperative child (7,10-13). If such patients are given intravenous free water (any solution with a sodium concentration below 140 mmol·l<sup>-1</sup>), there will always be a tendency towards postoperative hyponatraemia (14). When compared with other groups, prepubertal children are far more susceptible to brain damage from hyponatraemia than are adults (6), and recent experimental evidence demonstrates why this may be the case.

### *Effects of hyponatraemia on the paediatric central nervous system*

Nattie & Edwards (15) studied the effects of acute hyponatraemia on the brain of puppies. They found that acute lowering of plasma sodium from 140 to 120 mmol·l<sup>-1</sup> resulted in severe hypoxaemia (arterial PO<sub>2</sub> fell from 11.4-6.9 kPa (88 to 53 mmHg)) and cerebral oedema. In contrast to adults, the brains of paediatric animals (three day old puppies and neonatal rats) were unable to adapt to hypo-osmotic stress by extrusion of cation (15,16).

Adaptation of the brain to hyponatraemia occurs as a consequence of the following sequence of events. First, hyponatraemia leads to a movement of water into brain cells as a result of osmotic forces. In addition, vasopressin which is usually elevated in the plasma of hyponatraemic patients (17) may lead to a direct movement of water into brain cells independent of the effects of hyponatraemia (18). The early response of the brain to this hyponatraemia-mediated oedema is the loss of blood and cerebrospinal fluid, followed by extrusion of sodium from brain cells by several pathways (19). Loss of potassium and possibly organic osmolytes follows later, in an attempt to decrease brain cell osmolality without a gain of water (20).

### *Effects of hormones and physical factors on brain adaptation to hyponatraemia*

There is a significantly higher intracellular brain water content in prepubertal rats in comparison with adult rats, suggesting that the brain occupies a greater percent of the available intracranial volume in young rats (16). Such physical factors may be important determinants of outcome in hyponatraemic rats. As individuals age, there is a progressive decline in the volume of brain, while skull size remains constant in adult life (21). Thus, elderly individuals of both

Correspondence to: Allen L. Arieff, 299 South Street, Sausalito, CA 94965, USA.






Figure 1

In nine published series from our laboratory comprising 847 hospitalized patients with postoperative hyponatraemia, 19% (158/847) developed hyponatraemic encephalopathy and 117 developed permanent brain damage or died. The major risk factors associated with permanent brain damage in these 117 patients with hyponatraemic encephalopathy are shown. Most patients (96%) suffered an hypoxic episode because of failure to initiate active therapy in a timely manner. In 4% of patients suffering permanent brain damage, improper therapy for hyponatraemia was implicated in the outcome.

genders have more room in the rigid skull for the brain to expand than do younger ones. This finding is more marked in males (21).

If adaptation of the brain is not adequate, pressure of the swollen brain on the rigid skull leads to a decrease in cerebral blood flow (22) and cerebrospinal fluid production (23). If the ability of the brain to adapt is impaired, there will be increasing oedema, with eventual tentorial herniation and secondary cerebral ischaemia (24). This often leads to respiratory insufficiency (4), with reduced delivery of oxygen to brain because of the further decrease of cerebral blood flow, thereby exacerbating the existing cerebral ischaemia (22).

Sex steroid and certain neuropeptide hormones may influence brain adaptation to hyponatraemia. Male rabbits and cats are more efficient than females in extruding sodium to decrease brain cell osmolality during hyponatraemia, resulting in significantly less brain swelling in male than in female hyponatraemic animals (16,25). Oestrogens have also been reported to stimulate, and androgens to suppress, vasopressin release (26,27). Virtually all hyponatraemic patients have increased plasma levels of vasopressin (17,28), a neuropeptide which may exert multiple potentially deleterious cerebral effects. In normonatraemic animals vasopressin results in water accumulation in

the brain (18), a significant decline in brain synthesis of ATP (29), and a decline of brain pH (29,30). Vasopressin also impairs the function of several important adaptive pathways to hyponatraemia (31,32).

Recent studies have demonstrated that the brains of prepubertal rats are unable to adapt to hyponatraemia (16). The greater mortality with hyponatraemia in prepubertal rats is associated with a greater accumulation of water in the intracellular space of the brain than in rats belonging to other age groups, as well as an inability of the prepubertal brain to extrude sodium from brain cells. The baseline intracellular sodium content in the prepubertal rats was greater by almost 50% than in control adult rats, a finding consistent with previous studies in newborn dogs (15,33).

#### *Biochemical differences in paediatric vs adult brain with hyponatraemia*

There are several possible reasons for the increased brain intracellular sodium in prepubertal rats. The  $\text{Na}^+/\text{K}^+$  ATPase system appears to be the major early adaptive pathway for extrusion of sodium from brain cells during hyponatraemia (19,34) and its impairment results in decreased ability to pump sodium out of the brain. In prepubertal rats, the brain  $\text{Na}^+/\text{K}^+$  ATPase activity is significantly lower than that observed in adults, both *in vitro* (35) and *in vivo* (36). Coupled with the higher brain sodium, these differences may reflect a limited ability to pump sodium out of the prepubertal brain. The increased intracellular sodium content may be a consequence of limited cerebral  $\text{Na}^+/\text{K}^+$  ATPase function in young rats compared to adults. The decreased cerebral  $\text{Na}^+/\text{K}^+$  ATPase activity may be responsible for the impaired adaptation to hyponatraemia in prepubertal rats. Testosterone stimulates  $\text{Na}^+/\text{K}^+$  ATPase activity in rat brain (37,38). Pretreatment of prepubertal rats with testosterone resulted in a significant decrease in the brain intracellular content of both sodium and water while also reducing the mortality associated with acute hyponatraemia from 84% to zero (16).

#### *Clinical effects of hyponatraemia in children vs adults*

If one can extrapolate the above experimental findings to paediatric patients, then the implications would be that children are more susceptible to brain damage from postoperative hyponatraemia than are adults. The reasons include: a) decreased available



room 1 swelling of the paediatric brain in the rigid skull, leading to a propensity for brain herniation with what might appear to be a small decrement of plasma sodium (39); b) impaired ability of the paediatric brain to adapt to hyponatraemia when compared with adults (15,37); c) severe systemic hypoxaemia secondary to respiratory insufficiency frequently occurs in children with only modest hyponatraemia (6,15,39). The respiratory insufficiency is a consequence of increased intracranial pressure (3).

Gomola *et al.* have described a prepubertal (10 years old) female child with middle face hypoplasia who underwent elective maxillary reconstruction (40). The surgery went well and postoperatively, she was given primarily free water intravenously (280 mM glucose in 51 mM NaCl) at a rate of 2 l per day. The child weighed 30 kg with estimated total body water of 18.5 l. On the first postoperative day, the child became confused and developed headache and vomiting. Renal function was apparently normal on the basis of normal plasma urea and creatinine. The plasma sodium was found to be 117 mmol·l<sup>-1</sup>. She was initially treated with sodium supplementation, but on the second post-operative day, the plasma sodium was still low at 120 mmol·l<sup>-1</sup>. The urine and plasma osmolalities were 342 and 255 mOsm·kg<sup>-1</sup>. An MRI of the brain was normal. The authors proposed three possible explanations for the hyponatraemia: a) dilutional hyponatraemia secondary to IV hypotonic fluid; b) pituitary insufficiency; c) inappropriate secretion of ADH. Pituitary insufficiency was ruled out by normal values for ACTH, cortisol, thyroid hormone and growth hormone. The ADH was 4 to 5 pg·ml<sup>-1</sup>, which is 'normal' but inappropriately high for the extracellular hypoosmolality (41) and is essentially a universal finding in both paediatric and adult postoperative patients (7–13). The child received 2 l per day of hypotonic IV fluid in the presence of elevated plasma ADH. Although neither initial plasma sodium, urine output or total volume of IV fluids are provided, given the child's weight and rate of infusion, the plasma sodium of 117 mmol·l<sup>-1</sup> appears very likely to have been the consequence of retention of about 3 l of IV hypotonic fluid over two days (6). The expression inappropriate secretion of ADH (SIADH) was originally used for elevated plasma ADH related to lung cancer (42) and has become a catch all term for virtually any patient with elevated plasma ADH. In particular, postoperative patients as well as those with heart failure or hepatic

cirrhosis have elevated plasma ADH levels but are functionally hypovolaemic as well (41). Postoperative subjects are functionally hypovolaemic, so that the term SIADH may not be appropriate in this patient (11). There is also a perception that ADH, and by association SIADH, can somehow lower the plasma sodium. Although ADH leads to increased retention of ingested or infused water, in the absence of increased water intake, ADH by itself will have no effect upon the plasma sodium. Thus, the most likely explanation for the hyponatraemia in this patient is infusion of hypotonic fluid (51 mM NaCl/280 mM glucose) in the presence of the expected postoperative increase in plasma ADH. Adrenal insufficiency is ruled out by the normal plasma cortisol and the fact that she remained normal for six months without any steroid replacement therapy. Exactly why the plasma sodium rose following IV hydrocortisone is uncertain, but may have been related to the expected decline of ADH values to normal after four to five postoperative days. Pituitary insufficiency is ruled out by normal values for ACTH, IGF1 and growth hormone.

Symptomatic postoperative hyponatraemia carries a mortality of at least 15% (43), particularly in children and respiratory arrest is a frequent occurrence, but once this complication occurs, the morbidity is substantial (6,7). There is no obvious rationale for the administration of hypotonic fluid to a postoperative patient, unless the individual is hypernatraemic (14). If the patient becomes symptomatic, therapy with hypertonic NaCl is indicated (39). The syndrome can be prevented by administration of primarily isotonic fluids to postoperative patients.

## Acknowledgements

Supported by a grant RO1 AG 08575 from the National Institute on Aging, Department of Health and Human Services, Bethesda, MD, USA.

## References

- 1 Ayus JC, Arieff AI. Brain damage and postoperative hyponatremia: Role of gender. *Neurology* 1996; 46: 323–328.
- 2 Ayus JC, Wheeler JM, Arieff AI. Postoperative hyponatremic encephalopathy in menstruant women. *Ann Intern Med* 1992; 117: 891–897.
- 3 Ayus JC, Arieff AI. Pulmonary complications of hyponatremic encephalopathy: Noncardiogenic pulmonary edema and hypercapnic respiratory failure. *Chest* 1995; 107: 517–521.
- 4 Arieff AI. Hyponatremia, convulsions, respiratory arrest, and permanent brain damage after elective surgery in healthy women. *New Engl J Med* 1986; 314: 1529–1535.



- 5 Steele A, Gowrishankar M, Abrahamson S *et al*. Postoperative hyponatremia despite near-isotonic saline infusion: A phenomenon of desalination. *Ann Intern Med* 1997; 126: 20-25.
- 6 Arieff AI, Ayus JC, Fraser CL. Hyponatraemia and death or permanent brain damage in healthy children. *Brit Med J* 1992; 304: 1218-1222.
- 7 Cowley DM, Pabari M, Sinton TJ *et al*. Pathogenesis of postoperative hyponatraemia following correction of scoliosis in children. *Aust NZ J Surg* 1988; 58: 485-489.
- 8 Varavithya W, Hellerstein S. Acute symptomatic hyponatremia. *J Pediatrics* 1967; 71: 269-283.
- 9 Grushkin AB, Sarnaik A. Hyponatremia: Pathophysiology and treatment, a pediatric perspective. *Pediatr Nephrol* 1992; 6: 280-286.
- 10 Burrows FA, Shatack JG, Crone RK. Inappropriate secretion of antidiuretic hormone in a postsurgical pediatric population. *Crit Care Med* 1983; 11: 527-531.
- 11 Judd BA, Haycock GB, Dalton N *et al*. Hyponatraemia in premature babies and following surgery in older children. *Acta Paediatr Scand* 1987; 76: 385-393.
- 12 Judd BA, Haycock GB, Dalton N *et al*. Antidiuretic hormone following surgery in children. *Acta Paediatr Scand* 1990; 79: 461-466.
- 13 Brazel PW, McPhee IB. Inappropriate secretion of antidiuretic hormone in postoperative scoliosis patients: The role of fluid management. *Spine* 1996; 21: 724-727.
- 14 Rosenthal MH, Arieff AI. Fluid and electrolyte therapy in surgical and postoperative patients. *Curr Opinion Crit Care* 1995; 1: 469-477.
- 15 Nattie EE, Edwards WH. Brain and CSF water in newborn puppies during acute hypo- and hypernatremia. *J Appl Physiol* 1981; 51: 1086-1091.
- 16 Arieff AI, Kozniowska E, Roberts T *et al*. Age, gender and vasopressin affect survival and brain adaptation in rats with metabolic encephalopathy. *Am J Physiol* 1995; 268 (Reg Integr Comp Physiol 37) (5): R1143-R1152.
- 17 Anderson RJ, Chung HM, Kluge R *et al*. Hyponatremia: A prospective analysis of its epidemiology and the pathogenetic role of vasopressin. *Ann Intern Med* 1985; 102: 164-168.
- 18 Rosenberg GA, Estrada E, Kyner WT. Vasopressin-induced brain edema is mediated by the V1 receptor. *Adv Neurol* 1990; 52: 149-154.
- 19 Vexler ZS, Ayus JC, Roberts TPL *et al*. Ischemic or hypoxic hypoxia exacerbates brain injury associated with metabolic encephalopathy in laboratory animals. *J Clin Invest* 1994; 93: 256-264.
- 20 Melton JE, Patlak CS, Pettigrew KD *et al*. Volume regulatory loss of Na, Cl, and K from rat brain during acute hyponatremia. *Am J Physiol* 1987; 252 (Renal Fluid Electrolyte Physiol 21): F661-F669.
- 21 Gur RC, Mozley PD, Resnick SSM *et al*. Gender differences in age effect on brain atrophy measured by magnetic resonance imaging. *Proc Natl Acad Sci USA* 1991; 88: 2845-2849.
- 22 Kozniowska E, Roberts TPL, Vexler ZS *et al*. Hormonal dependence of the effects of metabolic encephalopathy on cerebral perfusion and oxygen utilization in the rat. *Circ Res* 1995; 76: 551-558.
- 23 Faraci FM, Mayhan WG, Heistad DD. Effect of vasopressin on production of cerebrospinal fluid: Possible role of vasopressin (V1)-receptors. *Amer J Physiol* 1990; 258 (Reg Integr Comp Physiol 27): R94-R98.
- 24 Fraser CL, Arieff AI. Fatal central diabetes mellitus and insipidus resulting from untreated hyponatremia: A new syndrome. *Ann Intern Med* 1990; 112: 113-119.
- 25 Vexler ZS, Roberts T, Derugin N *et al*. Adaptation of the brain to hyponatraemia in two different species: Effects of gender. *Abstracts Soc Neurosci* 1994; 24: (in press).
- 26 Akaishi T, Sakuma Y. Estrogen-induced modulation of hypothalamic osmoregulation in female rats. *Amer J Physiol* 1990; 258 (Reg Integr Comp Physiol 27): R924-R929.
- 27 Stone JD, Crofton JT, Share L. Sex differences in central adrenergic control of vasopressin release. *Amer J Physiol* 1989; 257: R1040-R1045.
- 28 Gross PA, Pehrish H, Rascher W *et al*. Pathogenesis of clinical hyponatremia: Observations of vasopressin and fluid intake in 100 hyponatremic medical patients. *Eur J Clin Invest* 1987; 17: 123-129.
- 29 Fraser CL, Kucharczyk J, Arieff AI *et al*. Sex differences result in increased morbidity from hyponatremia in female rats. *Am J Physiol* 1989; 256 (Reg Integr Comp Physiol 25): R880-R885.
- 30 Adler S, Simplaceanu V. Effect of acute hyponatremia on rat brain pH and rat brain buffering. *Amer J Physiol* 1989; 256 (Renal Electrol Physiol 25): F113-F119.
- 31 Kanda F, Arieff AI. Vasopressin inhibits calcium-coupled sodium efflux system in rat brain synaptosomes. *Amer J Physiol* 1994; 266 (Reg Integr Comp Physiol 35): R1169-R1173.
- 32 Kanda F, Sarnacki P, Arieff AI. Atrial natriuretic peptide inhibits the amiloride-sensitive sodium-hydrogen exchanger in rat brain. *Amer J Physiol* 1992; 263 (Reg Integr Comp Physiol 32): R279-R283.
- 33 Widdowson EM, Dickerson JWT. The effect of growth and function on the chemical composition of soft tissues. *Biochem J* 1960; 77: 30-43.
- 34 Fraser CL, Sarnacki P. Na<sup>+</sup>-K<sup>+</sup> ATPase pump function in male rat brain synaptosomes is different from that of females. *Am J Physiol* 1989; 257 (Endocr Metab 20): E284-E289.
- 35 Matsuda T, Shimizu I, Baba A. Postnatal change in a Ca<sup>2+</sup>-mediated decrease in (Na<sup>+</sup>+K<sup>+</sup>)-ATPase activity in rat brain slices. *Brain Res* 1992; 572: 349-351.
- 36 Jinna RR, Uzodinma JE, Desai D. Age-related changes in rat brain ATPases during treatment with chlordecone. *J Toxicol Environ Health* 1989; 27: 199-208.
- 37 Guerra M, del Castillo AR, Battaner E *et al*. Androgens stimulate preoptic area Na<sup>+</sup>, K<sup>+</sup> ATPase activity in male rats. *Neurosci Lett* 1987; 78: 97-100.
- 38 Fraser CL, Swanson RA. Female sex hormones inhibit volume regulation in rat brain astrocyte culture. *Am J Physiol (Cell Physiol 36)* 1994; 267: C909-C914.
- 39 Sarnaik AP, Meert K, Hackbarth R *et al*. Management of hyponatremic seizures in children with hypertonic saline: A safe and effective strategy. *Crit Care Med* 1991; 19: 758-762.
- 40 Gomola A, Cabrol S, Murat I. Severe hyponatraemia after plastic surgery in a girl with cleft palate, medial facial hypoplasia and growth retardation (case report). *Paediatr Anaesth* 1998; 8: 69-71.
- 41 Schrier RW. Pathogenesis of sodium and water retention in high-output and low-output cardiac failure, nephrotic syndrome, cirrhosis, and pregnancy. Parts 1 and 2. *New Engl J Med* 1988; 319(16): 1065-1072 and 1127-1134.
- 42 Bartter FE, Schwartz WB. The syndrome of inappropriate secretion of antidiuretic hormone. *Am J Med* 1967; 42: 790-806.
- 43 Fraser CL, Arieff AI. Epidemiology, pathophysiology and management of hyponatremic encephalopathy. *Am J Med* 1997; 102: 67-77.