

Ms Wendy Beggs Directorate of Legal Services 2 Franklin Street BELFAST BT2 8DQ Your Ref: NSC B04/1

Our Ref: CM-0028-11

Date: 29 September 2011

Dear Ms Beggs,

Re Investigation into the death of Adam Strain

I refer to Dr Taylor's Witness Statement Ref 008/3 dated 28/9/11 received yesterday by the Inquiry. The statement contains numerous references to documents without an Inquiry Reference number copies of which have not been provided with the statement. The Inquiry's statement request form clearly states that if a document is referred to by the witness without an Inquiry reference number then a copy of that document is to be provided to the Inquiry.

Please find attached a list of the documents referred to by Dr Taylor. A copy of these documents is required to be furnished by the Inquiry within 7 days from the date hereof.

Yours sincerely,

Caroline Martin

Solicitor to the Inquiry

Secretary: Bernie Conlon

Arthur House, 41 Arthur Street, Belfast, BT1 4GB

Emall: inquiry@ihrdni.org Website: www.ihrdni.org Tel: 028 9044 6340 Fax: 028 9044 6341

AS - INQ 301-068-454

Attachment

Cansick J, et al. A fatal case of cerebral oedema with hyponatraemia and massive polyuria after renal transplantation. Per O'Flynn PE and Milford CA. Fasting in children for day case surgery. Annals of the Royal College of Surgeons of England Thomas DKM. Hypoglycaemia in children before operation: Its incidence and prevention. BR J Anaesth 1974;46:66-8

Maze A, Samuels SI. Hypoglycaemia-induced seizures in an infant during anaesthesia. Anesthesiology 1980;52:77-8 Jenson BH, Wernberg M, Ansersen M. Pre-operative starvation and blood glucose concentrations in children undergoing in-patient and out-patient anaestheia. Br Anaesth 1982;54:1071-4

The Association of Anaesthetists of GB & Ireland, Monitoring Standards

Hatch and Sumner, Paediatric Anaesthesia 1999. P380 "superior vena cava pressures.... In the range of 10-15 mmHg" Mathur A, et al. Hypotonic vs isotonic saline solutions for intrevenous fluid management of acute infections. Cochrane Satabase Syst Rev. 2004;(2):CD004169)

Neville KA, et al. Isotonic is better than hypotonic saline for intravenous rehydration of children with gastroenteritis: a prospective randomised study. Arch Dis Child. 2006 Mar;91(3):226-32. Epub 2005 Dec 13)

Choong K, et al. Hyptonic versus isotonic saline in hospitalised children:a systematic review. Arch Dis Child. 2006 Oct;91(Yung M and Keeley S. Randomised controlled trial of intravenous maintenance fluids. J Paediatr Child Health. 2009 Jan-Feb;45(1-2):9-14 Epub 2007 Nov 25

Textbook of Pediatric Intensive Care, Rogers M 1987 p 656

Textbook of Pediatric Intensive Care, Rogers M 1987 p 627

AS - INQ 301-068-455